



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON A NEW TREATMENT OF THEOREMS OF FINITENESS*

BY

OLIVER EDMUNDS GLENN

Propositions relative to the finiteness of algebraic invariant systems can be treated under the theory of invariants of the type which involves the parameters of the transformations T to which a binary form

$$f_m = (a_0, \dots, a_m)(x_1, x_2)^m$$

is subjected. The algorithms of this theory are of extensive content. The full discussion of these functions comprehends not only the ordinary binary algebraic invariant theory as one of its phases, but, as was made clear in detail by the present author, in a former paper,[†] it comprehends, also, the general theory of orthogonal concomitants, of booleans, and of still other invariant systems appertaining to important special sets[‡] of binary substitutions. It might have been anticipated, I presume, that an invariant theory based upon considerations connected with the poles of the transformations, as the present theory is based, would be likely to be deeply grounded, but, prior to my own publication on this subject, no one seems to have adopted this point of view.

This paper contains a new and very simple proof of Gordan's theorem, and a determination of the complete system of concomitants in the domain $R(1, T, 0)$, of the binary quintic, carrying with it as one special case, for instance, the previously undetermined complete system of orthogonal concomitants of the quintic form.

As far as my knowledge goes, general proofs of the finiteness of orthogonal or of boolean systems have been given by no one, although these were among the first invariants discovered.[§] Apparently this was because researchers long took the erroneous view that these theories were fully included in that of simultaneous concomitants of the ordinary formal type. In this paper I prove these theorems for the general order m , and show that the systems are given by the finite set of irreducible solutions of a definite linear diophantine equation.

* Presented to the Society, December, 1917, and September, 1918.

† These Transactions, vol. 18 (1917), p. 443.

‡ Annals of Mathematics, vol. 20 (1918), p. 125.

§ Boole, Cambridge Mathematical Journal, vol. 3 (1843), p. 1.

1. CERTAIN FUNDAMENTAL SYSTEMS OF THE QUINTIC

The poles of the general transformation

$$T : \begin{cases} x_1 = \alpha_1 x'_1 + \alpha_2 x'_2, \\ x_2 = \beta_0 x'_1 + \beta_1 x'_2, \end{cases}$$

are the roots of the linear forms

$$(1_1) \quad f_{\pm 1} = 2\beta_0 x_1 + (\beta_1 - \alpha_1 \pm \Delta) x_2,$$

the latter being, accordingly, covariants of T . When f_m is expanded in terms of $f_{\pm 1}$ as arguments:

$$(1) \quad f_m = \sum_{i=0}^m \binom{m}{i} \phi_{m-2i} f_{+1}^{m-i} f_{-1}^i,$$

the coefficients ϕ of the expansion are linear invariants of f_m under T in the domain $R(1, T, \Delta)$, the notation for this domain indicating polynomials rational in the coefficients of T , in

$$\Delta = \sqrt{(\alpha_1 - \beta_1)^2 + 4\alpha_2 \beta_0},$$

and rational numerically. The invariant relations are

$$(2) \quad f'_{+1} = \rho_1^{-1} f_{+1}, \quad f'_{-1} = \rho_1^{-1} f_{-1},$$

where

$$\rho_{\pm 1} = \frac{1}{2} (\alpha_1 + \beta_1 \pm \Delta), \quad \rho_{+1} \rho_{-1} = D = \alpha_1 \beta_1 - \alpha_2 \beta_0;$$

and

$$(3) \quad \phi'_{m-2i} = \rho_1^{m-2i} D^i \phi_{m-2i} \quad (i = 0, \dots, m).$$

The invariants ϕ_{m-2i} can be given an explicit representation if a symbolical notation is adopted. Thus

$$f = a_x^m = (a_1 x_1 + a_2 x_2)^m = b_x^m.$$

and the inverse of the transformations

$$f_{\pm 1} = 2\beta_0 x_1 + (\beta_1 - \alpha_1 \pm \Delta) x_2,$$

is

$$T : \begin{cases} x_1 = (-4\beta_0 \Delta)^{-1} [(\gamma_1 - \Delta) f_{+1} - (\gamma_1 + \Delta) f_{-1}], \\ x_2 = (-4\beta_0 \Delta)^{-1} [2\beta_0 (-f_{+1} + f_{-1})] \end{cases} \quad (\gamma_1 = \beta_1 - \alpha_1).$$

Hence the expansion (1) is given by

$$f = (-4\beta_0 \Delta)^{-m} \{ (\gamma_1 - \Delta) a_1 - 2\beta_0 a_2 \} f_{+1} + \{ -(\gamma_1 + \Delta) a_1 + 2\beta_0 a_2 \} f_{-1} \}^m,$$

that is

$$(4) \quad \phi_{m-2i} = (-1)^i [(\gamma_1 - \Delta) a_1 - 2\beta_0 a_2]^{m-i} [(\gamma_1 + \Delta) a_1 - 2\beta_0 a_2]^i (-4\beta_0 \Delta)^{-m},$$

($i = 0, \dots, m$).

This notation will be especially useful in the second section of this paper.

If we differentiate ϕ_{m-2i} according to the operator

$$a_1 \frac{\partial}{\partial a_2} + 2\beta_0 \frac{\partial}{\partial \gamma_1} - \gamma_1 \frac{\partial}{\partial \alpha_2},$$

the result is found to vanish. Hence, since

$$a_1 \frac{\partial}{\partial a_2} = a_1 \left[\frac{\partial}{\partial a_0} \frac{\partial a_0}{\partial a_2} + \frac{\partial}{\partial a_1} \frac{\partial a_1}{\partial a_2} + \cdots + \frac{\partial}{\partial a_m} \frac{\partial a_m}{\partial a_2} \right],$$

the invariant ϕ_{m-2i} satisfies the differential equation

$$\left(a_0 \frac{\partial}{\partial a_1} + 2a_1 \frac{\partial}{\partial a_2} + \cdots + ma_{m-1} \frac{\partial}{\partial a_m} + 2\beta_0 \frac{\partial}{\partial \gamma_1} - \gamma_1 \frac{\partial}{\partial \alpha_2} \right) \phi_{m-2i} = 0.$$

Every invariantive function, rational or irrational, of the coefficients and variables of f_m , of the coefficients of T , and of Δ , is expressible in terms of the ϕ -s and of $f_{\pm 1}$. We have only to employ the inverse of the set of linear equations (in a_0, \cdots, a_m)

$$\phi_{m-2i} = \phi_{m-2i}(a_0, \cdots, a_m) \quad (i = 0, \cdots, m).$$

In the domain $R(1, T, \Delta)$ a complete system is composed of ϕ_{m-2i} ($i = 0, \cdots, m$), f_{+1} , f_{-1} .

In the domain $R(1, T, 0)$, rational in the coefficients of T , a complete system is in one-to-one correspondence with those invariantive products

$$(5) \quad P = \prod_{i=0}^m \phi_{m-2i}^{x_i} f_{+1}^{\sigma_1} f_{-1}^{\sigma_2}$$

for which the exponent of ρ_1 in the invariant relation

$$P' = \rho_1^a D^b P$$

is zero, and for this reason complete systems in this domain are given by the complete set of irreducible solutions in positive integers of the linear diophantine equation

$$(6) \quad a = \prod_{i=0}^m (m - 2i) x_i - \sigma_1 + \sigma_2 = 0.$$

The particularization of T which changes the systems belonging to the domain $R(1, T, 0)$ into boolean concomitants is given by

$$\alpha_1 = \frac{\sin(\omega - \alpha)}{\sin \omega}, \quad \alpha_2 = \frac{\sin(\omega - \beta)}{\sin \omega}, \quad \beta_0 = \frac{\sin \alpha}{\sin \omega}, \quad \beta_1 = \frac{\sin \beta}{\sin \omega}.$$

When $\omega = \frac{1}{2}\pi$ the latter systems become orthogonal. We have now proved the following

THEOREM. *Complete systems in the domain $R(1, T, 0)$, booleans, ortho-*

gonal systems, as well as concomitants of f_m under the so-called relativity* transformations

$$T : \begin{cases} t = \frac{1}{\sqrt{1 - \beta^2}} \left(t' + \frac{v}{c^2} x' \right), \\ x = \frac{1}{\sqrt{1 - \beta^2}} (x' + vt'), \end{cases}$$

are finite, and are given by the finite set of irreducible solutions in positive integers of the equation (6).

In the instance $m = 5$, therefore, we have to solve the equation

$$(7) \quad 5x_0 + 3x_1 + x_2 + \sigma_2 = \sigma_1 + x_3 + 3x_4 + 5x_5$$

and if an irreducible solution is given by the ξ -s in the first row in the table following, another irreducible solution representing the product conjugate to the first product is furnished by the second row:

x_0	x_1	x_2	σ_2	x_3	x_4	x_5	σ_1
ξ_0	ξ_1	ξ_2	s_2	ξ_3	ξ_4	ξ_5	s_1
ξ_5	ξ_4	ξ_3	s_1	ξ_2	ξ_1	ξ_0	s_2

If the two products P, Q thus found are combined, giving $P \pm Q = \alpha_{\pm 1}$, then the $\alpha_{\pm 1}$, deprived of certain irrelevant constant factors, are members of the fundamental system. I have carried out the work of solving (7) and I give in the table below the set of irreducible solutions found. For convenience in reading off the concomitants $P \pm Q$ a solution and its conjugate are juxtaposed under a double notation, as $l_{\pm 1}$, the concomitants $l_{\pm 1}$ being, evidently,

$$(8) \quad l_{\pm 1} = \phi_3 \phi_{-1}^2 f_{+1} \pm \phi_{-3} \phi_1^2 f_{-1}.$$

The number of invariants and covariants in this fundamental system including $f_{+1}f_{-1}$, is 42, of which 15 are pure invariants. The solutions furnishing the latter are placed in the top rows of the table.

A few remarks should perhaps be added as to the method of solving (7). Transposition of terms reduces the equation to

$$(9) \quad 5x + 3y + z = 0,$$

where

$$(9_1) \quad x_0 - x_5 = x, \quad x_1 - x_4 = y, \quad x_2 - x_3 - \sigma_1 + \sigma_2 = z.$$

The equation (9) is now solved for its irreducible solutions in both positive and negative integers x, y, z , and to each of these solutions correspond sets for (7), given by (9₁). The problem is thus divided into a large number of simple mutually exclusive sub-problems.

* R. D. Carmichael, *Theory of Relativity*, p. 44.

	x_0	x_1	x_2	x_3	x_4	x_5	σ_1	σ_2
$a \dots$	1					1		
$b \dots$		1			1			
$c \dots$			1	1				
$d_{\pm 1} \dots \{$		1	3	3	1			
$e_{\pm 1} \dots \{$	1		5	5		1		
$g_{\pm 1} \dots \{$	1	1	2	2	1	1		
$h_{\pm 1} \dots \{$	1	2	1	1	2	1		
$i_{\pm 1} \dots \{$	3	5			5	3		
$j_{\pm 1} \dots \{$	2	3	1	1	3	2		
$k_{\pm 1} \dots \{$			1	1			1	1
$l_{\pm 1} \dots \{$		1	2	2	1		1	1
$m_{\pm 1} \dots \{$		1	1	1	1		2	2
$n_{\pm 1} \dots \{$		1			1		3	3
$o_{\pm 1} \dots \{$	1		4	4		1	1	1
$p_{\pm 1} \dots \{$	1		3	3		1	2	2
$q_{\pm 1} \dots \{$	1		2	2		1	3	3
$r_{\pm 1} \dots \{$	1		1	1		1	4	4
$s_{\pm 1} \dots \{$	1					1	5	5
$t_{\pm 1} \dots \{$	1	1	1	1	1	1	1	1
$u_{\pm 1} \dots \{$	1	1			1	1	2	2
$v_{\pm 1} \dots \{$	1	2			2	1	1	1
$w_{\pm 1} \dots \{$	2	3			3	2	1	1

2. THE FINITENESS OF ALGEBRAIC SYSTEMS

Consider now the following well-known lemma on systems of polynomials, first proved by Hilbert: If an infinite system of homogeneous forms in n variables satisfies any law sufficiently explicit to locate an arbitrarily chosen homogeneous form within or without the system, then there always exists a set of these forms, finite in number, as F_1, \dots, F_s , each within the system,

and such that any form F belonging to the system is expressible in the form

$$(10) \quad F = T_1 F_1 + \cdots + T_s F_s,$$

where T_1, \cdots, T_s are also homogeneous polynomials in the same n variables but do not necessarily belong to the original system.*

The generality of this theorem with regard to the law or laws which determine the system of forms under consideration makes it applicable to a variety of problems, but the possibilities in connection with the study of particular given laws of formation and the systems appertaining have not been much exploited in literature.

When we apply the methods of this paper to systems of concomitants in the domain $R(1, 0, 0)$, functions which are free from the coefficients of the transformation T and therefore are the ordinary rational integral concomitants of f_m , we discover a very interesting application of this lemma. The character of (1) as a typical representation of f_m immediately shows that any concomitant C of f_m in $R(1, 0, 0)$ is a homogeneous polynomial F in the invariants ϕ_{m-2i} ($i = 0, \cdots, m$), f_{+1}, f_{-1} , and formed according to a law sufficiently definite to define a system of polynomials in the technical sense of the lemma. This law is that F shall be such a polynomial that, although its variables $\phi_{m-2i}, f_{+1}, f_{-1}$ belong to the domain $R(1, T, \Delta)$, it simplifies by multiplying out into an expression (in $x_1, x_2, a_0, \cdots, a_m$) belonging to the domain $R(1, 0, 0)$, the parameters $\alpha_1, \alpha_2, \beta_0, \beta_1$ disappearing by cancellation from numerator and denominator.

Illustration. If $m = 2$ we have; ($\gamma_1 = \beta_1 - \alpha_1$),

$$\phi_{\pm 2} = (2^3 \beta_0^2 \Delta^2)^{-1} [(\gamma_1^2 + 2\alpha_2 \beta_0 \mp \gamma_1 \Delta) a_0 + 2\beta_0 (-\gamma_1 \pm \Delta) a_1 + 2\beta_0^2 a_2],$$

$$\phi_0 = (2\beta_0 \Delta^2)^{-1} [\alpha_2 a_0 + \gamma_1 a_1 - \beta_0 a_2],$$

and

$$4\beta_0^2 \Delta^2 (\phi_0^2 - 4\phi_2 \phi_{-2}) \equiv a_1^2 - a_0 a_2 \quad (= F).$$

We can now prove that the system of concomitants belonging to the domain $R(1, 0, 0)$ is finite (Gordan's theorem). For, suppose that F is any such concomitant. It is a homogeneous polynomial in ϕ_{m-2i} ($i = 0, \cdots, m$), f_{+1}, f_{-1} , as above. Hence there exists a finite set of polynomials in $\phi_{m-2i}, f_{+1}, f_{-1}$, as F_1, \cdots, F_s , each simplifying by multiplication into a concomitant belonging to the domain $R(1, 0, 0)$, such that

$$(10) \quad F = T_1 F_1 + \cdots + T_s F_s,$$

and the polynomials T_i , being homogeneous in the invariants $\phi_{m-2i}, f_{\pm 1}$, are concomitants which may or may not belong to the domain $R(1, 0, 0)$.

* *Mathematische Annalen*, vol. 36 (1890), p. 473.

We can now prove that the T -s do, however, belong, in fact, to this domain. This will be equivalent to a proof of the finiteness theorem as each T function can then be expressed linearly in terms of F_1, \dots, F_s and the process repeated until F is reduced to a polynomial in the forms of the finite fundamental system F_1, \dots, F_s .

The following propositions should now be noted: (a) The selection of the set of concomitants F_i in (10) is not unique. The expression of F as a polynomial in $f_{+1}, f_{-1}, \phi_{m-2i}$ ($i = 0, \dots, m$) is, on the other hand, unique, since we can obtain this expression (eq. (10)) by the substitution for $x_1, x_2, a_0, \dots, a_m$ in F , the linear functions τ in f_1, f_{-1} , and the linear functions in ϕ_m, \dots, ϕ_{-m} obtained by solving for the inverse of equations (4). Note that the equations (4) are consistent for this solution. If the determinant of this linear system were zero, the ϕ_{m-2i} would be linearly connected; then a_0, \dots, a_m would be linearly dependent. But these are arbitrary.

Although F is a unique polynomial P in $f_1, f_{-1}, \phi_m, \dots, \phi_{-m}$ and (10) is merely a grouping of the terms of P into components such that F_1, \dots, F_s are concomitants which simplify into the domain $R(1, 0, 0)$, nevertheless, even after the non-unique set F_1, \dots, F_s is definitely selected, the set of auxiliary factors T_1, \dots, T_s is not necessarily unique.*

(b) The linear transformation T is generated by combination of the following particular transformations:

$$\begin{aligned} T_n : x_1 &= \xi + n\eta, & x_2 &= \eta; \\ S_k : x_1 &= \xi, & x_2 &= k\eta \ (k \neq 0); \\ V : x_1 &= -\eta, & x_2 &= \xi. \end{aligned}$$

Invariancy under V is equivalent to the property of symmetry of invariants under the permutations $(a_0 a_m)(a_1 a_{m-1}) \dots (x_1 x_2)$. The transformation S_k represents isobarism, and T_n , seminvariance. It follows that a homogeneous polynomial in $a_0, \dots, a_m, x_1, x_2$ which possesses the properties of symmetry, isobarism, and seminvariance is a concomitant under T , and conversely.

(c) The equations of τ , looked upon as linear transformations are as general as those of T .

(d) If f is transformed into $f' = (a'_0, \dots, a'_m | x_1, x_2)^m$ by a general linear transformation t and if $I' = I(a'_0, \dots, a'_m; x'_1, x'_2)$ is any homogeneous polynomial in its arguments which possesses the technical properties of symmetry, isobarism, and seminvariance, then I' is the left-hand member of the invariant relation of an invariant I , viz.,

$$(11) \quad I(a'_0, \dots; x'_1, x'_2) = D^\Delta I(a_0, \dots; x_1, x_2).$$

* Cf. some theory by the present writer, on expressing a polynomial in terms of others, in these Transactions, vol. 15 (1914), p. 77, and vol. 18 (1917), p. 443.

To prove this, notice that the assumed properties of symmetry, isobarism, and seminvariance are sufficient conditions for the invariance of I' under the transformation of f' by a general linear transformation t' . But if f' has a formal invariant I' , under t' , then, by correspondence, f has a formal invariant $I(a_0, \dots, a_m; x_1, x_2)$ under t . Hence I' cannot be other than the transformed function of $I(a_0, \dots, a_m; x_1, x_2)$ under transformation of f by t , which was to be proved.

(e) The concomitants T_i belong to the domain $R(1, T, 0)$ if not to $R(1, 0, 0)$. For, if f' is the transformed of f by T , the relation for f' analogous to (10) is of the type

$$(\alpha\beta)^k \cdot F = T'_1(\alpha\beta)^{k_1} F_1 + \dots + T'_s(\alpha\beta)^{k_s} F_s \quad [(\alpha\beta) = D],$$

and this relation is necessarily identical with (10). Thus the invariant relation for T_i is of the form

$$T'_i = (\alpha\beta)^{k'_i} T_i,$$

where k'_i is a positive integer. Since the multiplier in this invariant relation involves D , and neither of its irrational factors ρ_1, ρ_{-1} , this amounts to a sufficient condition that T_i belong to $R(1, T, 0)$.

Consider next the equation (1) of Section 1. This form $f' = f_m$ is the result of transforming f_m by a binary substitution τ of general type (cf. proposition (c)). We wish to show that the auxiliary factors T_1, \dots, T_s in (10) are functions I' of proposition (d), i. e., they are functions of ϕ_{m-2i} ($i = 0, \dots, m$), f_{+1}, f_{-1} which possess the formal properties of symmetry, isobarism, and seminvariance.

(A) *Symmetry*. That T_i is unaltered, save possibly by sign, by the permutational substitution $S = (\phi_m \phi_{-m}) (\phi_{m-2} \phi_{-m+2}) \dots (f_1 f_{-1})$ follows immediately from the fact that T_i belongs to the domain $R(1, T, 0)$. For, it must be a linear function of combinations $P + Q$, or of $P - Q$, where Q is a product of powers of ϕ_{m-2i} ($i = 0, \dots, m$), f_{+1}, f_{-1} and P is the product conjugate to Q . But the substitution S interchanges P and Q (cf. (8)).

(B) *Isobarism*. The isobaric property of T_i implies, in this instance, its invariance under the transformations

$$(12) \quad \phi'_{m-2i} = k^i \phi_{m-2i}; \quad f'_{+1} = f_{+1}, \quad f'_{-1} = k^{-1} f_{-1}$$

(cf. proposition (b)).

Now a representative term of T_i is (cf. (5))

$$P = \phi_m^{x_0} \phi_{m-2}^{x_1} \phi_{m-4}^{x_2} \dots f_{+1}^{\sigma_1} f_{-1}^{\sigma_2},$$

and both $\sigma_1 + \sigma_2$ and $x_0 + x_1 + \dots + x_m$ are invariable. If the transformations (12) are applied to P the exponent of the power of k which is a factor of the result is

$$w = x_1 + 2x_2 + \dots + mx_m - \sigma_2.$$

But from equation (6),

$$w = \frac{1}{2} [m(x_0 + x_1 + \cdots + x_m) - (\sigma_1 + \sigma_2)],$$

and this number is evidently invariable for the various terms P of T_i . Hence the isobaric property is proved.

(C) *Seminvariance.* The seminvariance of T_i is the property of remaining absolutely unaltered when we apply to it those substitutions upon ϕ_{m-2i} ($i = 0, \dots, m$), f_{+1} , f_{-1} which are induced by the transformations T_n of proposition (b), viz.,

$$(13) \quad f_{+1} = f'_{+1} + n f'_{-1}, \quad f_{-1} = f'_{-1},$$

applied to f_m of (1).

Let us suppose now that n is infinitesimal. Then, by a well-known bit of elementary analysis which need not be repeated here, the result of transforming T_i by the induced transformations is

$$(14) \quad T'_i = T_i + n \left[\phi_m \frac{\partial}{\partial \phi_{m-2}} + 2\phi_{m-2} \frac{\partial}{\partial \phi_{m-4}} + \cdots \right. \\ \left. + m\phi_{-m+2} \frac{\partial}{\partial \phi_{-m}} - f_{-1} \frac{\partial}{\partial f_{+1}} \right] T_i.$$

Moreover, when the induced transformations are applied to both members of equation (10), F and F_i ($i = 1, \dots, s$) remain unaltered since these functions reduce to ordinary covariants in the domain $R(1, 0, 0)$. Then this result of transforming (10) takes the form

$$F = T'_1 F_1 + T'_2 F_2 + \cdots + T'_s F_s.$$

Therefore, unless the quantity in brackets in (14) vanishes, this expression for F is one of the other possible representations of F as a linear function of F_i ($i = 1, \dots, s$) constructed upon the principle that the set of auxiliary factors T_i is not unique (cf. proposition (a)). But this is impossible by (B), for, unless the bracket in (14) is zero, T'_i is not isobaric. Hence $T'_i = T_i$ and the property of seminvariance is established.

Conclusion. Thus, by the principle of correspondence, in accordance with proposition (d), we have established the relations

$$(15) \quad T_i(\phi_m, \phi_{m-2}, \dots; f_{+1}, f_{-1}) = R^\lambda T_i(a_0, a_1, \dots; x_1, x_2) \\ (i = 1, \dots, s),$$

where R is the determinant of τ . We can now prove that $\lambda = 0$, so that T_i is absolutely equal to a concomitant belonging to the domain $R(1, 0, 0)$. This is done by showing that T_i is of degree zero in the parameters $\alpha_1, \alpha_2, \beta_0, \beta_1$, a fact which is fairly evident from empirical considerations, but an analytic proof of which is not without interest.

Note that the T -s, though fractional in the parameters are homogeneous in them, and hence, by Euler's theorem,

$$\nabla T_i = \left(\alpha_1 \frac{\partial}{\partial \alpha_1} + \alpha_2 \frac{\partial}{\partial \alpha_2} + \beta_0 \frac{\partial}{\partial \beta_0} + \beta_1 \frac{\partial}{\partial \beta_1} \right) T_i = t'_i T_i,$$

where t'_i is the degree of T_i in the parameters. Now suppose that

$$t'_1 = t'_2 = \cdots = t'_r$$

and operate by ∇ upon both members of (10). Since the F -s are free from α_1 , α_2 , β_0 , β_1 , the result is

$$t'_1 (T_1 F_1 + \cdots + T_r F_r) + \cdots + t'_s F_s = 0;$$

or

$$(16) \quad t_1 P_1 + t_2 P_2 + \cdots + T_v P_v = 0,$$

where the degrees t_1, \cdots, t_v are all different and $v \geq 1$. Next operate $v - 1$ times in succession by ∇ upon (16), and we get the linear system

$$t_1^k P_1 + t_2^k P_2 + \cdots + t_v^k P_v = 0 \quad (k = 1, \cdots, v).$$

The determinant C of this linear system is the circulant of the v distinct numbers t_i , i. e.,

$$C = \begin{vmatrix} t_1 & t_2 & \cdots & t_v \\ t_1^2 & t_2^2 & \cdots & t_v^2 \\ \vdots & \vdots & \ddots & \vdots \\ t_1^v & t_2^v & \cdots & t_v^v \end{vmatrix} \neq 0.$$

We would therefore have all P -s zero, and hence the contradiction $F = 0$, against which eventuality there are only two alternatives, viz., first, that the t -s should all be zero due to the fact that the numerator and denominator of each function T_i are of the same degree in the parameters, or, secondly, that the T -s are free from the parameters. Thus $\lambda = 0$, and (15) shows that the T_i are concomitants belonging to $R(1, 0, 0)$. This completes the proof of the theorem of finiteness.

UNIVERSITY OF PENNSYLVANIA